

# NEW STANDARD ACADEMY

Marks: 60

Date : 21-05-24

CLASS : 12<sup>TH</sup> (NEET)

Time: 90 min.

## PHYSICS

1. Four electric charge  $+q, +q, -q$  and  $-q$  are placed at the corners of a square of side  $2L$  the electric potential at a point A, Midway between two charges  $+q$  and  $+q$ , is

- a)  $\frac{1}{4\pi\epsilon_0} \frac{2q}{L} \left[ 1 + \frac{1}{\sqrt{5}} \right]$   
 b)  $\frac{1}{4\pi\epsilon_0} \frac{2q}{L} \left[ 1 - \frac{1}{\sqrt{5}} \right]$   
 c) Zero  
 d)  $\frac{1}{4\pi\epsilon_0} \frac{2q}{L} \left[ 1 + \sqrt{5} \right]$

2. Four point charges  $-Q, -q, 2q$  and  $+2Q$  placed one at each corner of the square. the relation between  $Q$  and  $q$  for which potential at the centre of square is zero is

- a)  $Q = -q$       b)  $Q = -\frac{1}{q}$   
 c)  $Q = q$       d)  $Q = \frac{1}{q}$

3. A capacitor having capacity of  $2\mu F$  is charged to 200V and then the plates of the capacitor are connected to a resistance wire. The heat produced in joule will be

- a)  $2 \times 10^{-2}$       b)  $4 \times 10^{-2}$   
 c)  $4 \times 10^4$       d)  $4 \times 10^{10}$

4. A capacitor is charged to 200 volt has 0.1 Coulomb charge. When it is discharged energy will be

- a) 1J      b) 4 J  
 c) 10 J      d) 20J

5. If a dielectric substance is introduced between the plates of a charged air-gap capacitor. The energy of the capacitor will
- a) Increase      b) Decrease  
 c) Remain unchanged  
 d) First decrease and then increase


6. The potentials of the two plates of capacitor are  $+10V$  and  $-10V$ . The charge on one of the

plates is  $40\mu C$ . The capacitance of the capacitor is


- a) 2F      b) 4F  
 c) 0.5F      d) 0.25F

7. The charge on  $4\mu F$  capacitor in the given circuit is.... in  $\mu C$ .

- a) 12  
 b) 24  
 c) 36  
 d) 32




8. Effective capacitance between A and B in the figure shown is (all capacitance are in  $\mu F$ )



- a)  $21\mu F$       b)  $23\mu F$   
 c)  $\frac{3}{14}\mu F$       d)  $\frac{14}{3}\mu F$

9. In the figure shown the equivalent capacitance between A and B is:

- a) 3.75F  
 b) 2F  
 c) 21F  
 d) 16F



10. When a slab of dielectric material is introduced between the parallel plates of a capacitor which remains connected to a battery

, Then charge on plates relative to earlier charge

- a) Is less
- b) Is same
- c) Is more
- d) May be less or more depending on the nature of the material introduced

### CHEMISTRY

11. For the chemical reaction  $3O_2 \rightarrow 2O_3$  the rate of formation of  $O_3$  is  $0.04 \text{ mole L}^{-1} \text{ sec}^{-1}$ . determine the rate of disappearance of  $O_2$ .

- a)  $0.04 \text{ mole L}^{-1} \text{ sec}^{-1}$
- b)  $0.08 \text{ mole L}^{-1} \text{ sec}^{-1}$
- c)  $0.10 \text{ mole L}^{-1} \text{ sec}^{-1}$
- d)  $0.06 \text{ mole L}^{-1} \text{ sec}^{-1}$

12. The values of rate constant for the decomposition of  $N_2O_5$ ,  $N_2O_5 \rightarrow N_2O_4 + \frac{1}{2}O_2$  are  $3.50 \times 10^{-5}$  and  $5 \times 10^{-3}$  at  $27^\circ\text{C}$  and  $67^\circ\text{C}$ , respectively. calculate the energy of activation?

- a) 14.8 kcal/mol
- b) 24.8 kcal/mol
- c) 25.31 kcal/mol
- d) 34.8 kcal/mol


13. The rate of reaction increases by 2.3 times when the temperature is raised from  $300\text{K}$  to  $310\text{K}$ . If  $K$  is the rate constant at  $300\text{K}$  then the rate constant at  $310\text{K}$  will be equal to

- a)  $2K$
- b)  $K$
- c)  $2.3K$
- d)  $3K^2$



14. In a first order reaction  $A \rightarrow \text{Products}$ , the ratio of  $a$  and  $(a-x)$  was found to be 8 after 60 minutes. Calculate the rate of the reaction in moles of  $A$  reacted per minute, if its concentration is  $0.1 \text{ mol L}^{-1}$

- a)  $5.566 \times 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1}$
- b)  $3.466 \times 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1}$
- c)  $4.366 \times 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1}$
- d)  $3.466 \times 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1}$

15. The reaction



Follows the mechanism given below



The overall order of this reaction is

- a) 2
- b) 1
- c) 3
- d) 0

16. In the following first order completing reactions.  $A + \text{Reagent} \rightarrow \text{Product}$ ,  $B + \text{Reagent} \rightarrow \text{Product}$ , the ratio of  $K_1/K_2$  If only 59% of  $B$  will have been reacted, When 94% of  $A$  has been reacted is

- a) 4.06
- b) 0.246
- c) 2.06
- d) 0.06

17. Which of the following is not an anodic reaction?

- a)  $Ag^+ \rightarrow Ag + e^-$
- b)  $Cu \rightarrow Cu^{2+} + 2e^-$
- c)  $Fe^{2+} \rightarrow Fe^{3+} + e^-$
- d)  $4OH^- \rightarrow 2H_2O + O_2 + 4e^-$

18. The standard e.m.f of a cell, involving one electron change is found to be  $0.591\text{V}$  at  $25^\circ\text{C}$ . The equilibrium constant of the reaction is ( $F=96500 \text{ C mol}^{-1}$ ;  $R=8.314 \text{ JK}^{-1} \text{ mol}^{-1}$ )

- a)  $1.0 \times 10^{10}$
- b)  $1.0 \times 10^5$
- c)  $1.0 \times 10^1$
- d)  $1.0 \times 10^{30}$

19. Calculate the quantity of electricity that would be required to reduce  $12.3\text{g}$  of nitrobenzene to aniline if current efficiency is 50%. If the potential drops across the cell is  $3.0\text{volts}$  how much energy will be consumed?

- a) 347.4KJ
- b) 447.4 KJ
- c) 3474 KJ
- d) 3.474KJ

20. If  $v$ , in the equation  $\Lambda = sp. cond \times V$ , is the volume in cc containing 1 eq. of the electrolyte;  $V$  for a  $\frac{N}{10}$  solution will be

- a) 10 cc
- b) 100cc
- c) 1000 cc
- d) 10000 cc

### BIOLOGY

21. Match the list of items of column I with column II and select the correct option from the codes given below:

column I	column II
A. F.Meischer	i) DNA duble helix
B. Griffith	ii) Nuclein
C. Hershey and Chase	iii) S.Pneumoniae
D. Watson and Crick	iv) Bacteriophages
E. Wilkins and Franklin	v) X-ray diffraction

Studies

- a) A-(ii),B-(iii),C-(iv),D-(i),E-(v)
- b) A-(ii),B-(iv),C-(iii),D-(i),E-(v)
- c) A-(i),B-(iii),C-(iv),D-(ii),E-(v)

- d) A-(i),B-(iv),C-(iii),D-(ii),E-(v)
22. Histone proteins are
- Basic, negatively charged
  - Basic, Positively charged
  - Acidic Positively charged
  - Acidic ,negatively charged
23. DNA dependent RNA polymerase catalyzes the polymerization in
- 5'-3' direction
  - 3'-5' direction
  - 3'-1' direction
  - 1'-3' direction
24. The sequence of structural genes in lac operonis
- Lac A, Lac Y, Lac Z
  - Lac A, Lac Z, Lac Y
  - Lac Y, Lac Z, Lac A
  - Lac Z, Lac Y, Lac A
25. The probes used in DNA fingerprinting technique are
- Radioactive natural DNA/RNA with known sequences
  - Radioactive synthetic DNA/RNA with unknown sequences
  - Radioactive natural DNA/RNA with unknown sequences
  - Radioactive synthetic DNA/RNA with known sequences
26. Which of the following sequence of steps is correct in DNA fingerprinting?
- Southern blotting, Electrophoresis, Hybridization, Autoradiography
  - Autoradiography, Electrophoresis, Hybridization, Southern blotting
  - Electrophoresis, Southern blotting, Hybridization, Autoradiography
  - Hybridization, Southern blotting, Electrophoresis, Autoradiography
27. Hypervariable region of DNA is formed of
- Minisatellite DNA
  - Microsatellite DNA
  - Probes
  - Both (a) and (b)
28. What is the criterion for DNA Fragments movement on agarose gel during gel electrophoresis?
- The larger the fragment size, the the farther it moves.
  - The smaller the fragment size, the the farther it moves.

- Positively charged fragments move to the father end.
- Negatively charged fragments donot move.

The question given below consists of Assertion and Reason. Use the following key to select the correct answer:

- If both assertion and reason are correct and reason is correct explanation for assertion.
  - If both assertion and reason are correct but reason is not correct explanation for assertion.
  - IF assertion is correct but reason is incorrect.
  - Both assertion and reason is incorrect.
29. **Assertion(A):** Split genes concept is applicable only to the prokaryotes.  
**Reason(R):** Prokaryotic genome is divided into exons and introns
30. **Assertion(A):** Replication and transcription occur in the nucleus but translation occurs in the cytoplasm.  
**Reason:** mRNA is transferred from the the nucleus into the cytoplsm where ribosomes and amino acids are available for protein synthesis